



August Forecast Update for Northwest Pacific Typhoon Activity in 2005

Issued: 5th August 2005

by Dr Adam Lea and Professor Mark Saunders
Benfield Hazard Research Centre, UCL (University College London), UK

Forecast Summary

TSR continues to anticipate the 2005 Northwest Pacific typhoon season will see activity close to average.

The TSR (Tropical Storm Risk) August forecast update for Northwest Pacific typhoon activity in 2005 continues to anticipate a near-average activity season. The forecast spans the full Northwest Pacific season from 1st January to 31st December 2005 (95% of typhoons historically occur after 1st May) and is based on data available through the end of July 2005. The forecast includes deterministic and probabilistic projections for overall basin activity, and deterministic projections for the numbers of tropical storms, typhoons and intense typhoons. TSR anticipates activity will be slightly above average for tropical storm numbers, typhoon numbers, intense typhoon numbers and the ACE index. TSR's main predictor at this lead for overall activity is the forecast anomaly in August-September Niño 3.75 sea surface temperature (SST) which we anticipate will be $0.28 \pm 0.20^\circ\text{C}$ warmer than normal this summer.

NW Pacific ACE Index and System Numbers in 2005

		ACE Index	Intense Typhoons	Typhoons	Tropical Storms
TSR Forecast (\pm FE)	2005	328 (\pm 78)	9.4 (\pm 2.5)	17.5 (\pm 2.9)	27.6 (\pm 3.7)
40yr Climate Norm (\pm SD)	1965-2004	305 (\pm 99)	8.6 (\pm 3.0)	16.9 (\pm 3.7)	26.8 (\pm 4.5)
Forecast Skill at this Lead	1965-2004	38%	33%	37%	33%

- Key: ACE Index = Accumulated Cyclone Energy Index = Sum of the Squares of 6-hourly Maximum Sustained Wind Speeds (in units of knots) for all Systems while they are at least Tropical Storm Strength. ACE Unit = $\times 10^4$ knots².
- Intense Typhoon = 1 Minute Sustained Wind > 95Kts = Hurricane Category 3 to 5
- Typhoon = 1 Minute Sustained Wind > 63Kts = Hurricane Category 1 to 5
- Tropical Storm = 1 Minute Sustained Wind > 33Kts
- SD = Standard Deviation
- FE (Forecast Error) = Standard Deviation of Errors in Simulated Real Time Forecasts 1965-2004
- Forecast Skill = Percentage Reduction in Mean Square Error Afforded by Cross-Validated Hindcasts 1965-2004 over Hindcasts Made with the 1965-2004 Climate Norm.
- Northwest Pacific = Northern Hemisphere Region West of 180°W Including the South China Sea. Any Tropical Cyclone (Irrespective of Where it Forms) Which Reaches Tropical Storm Strength Within this Region Counts as an Event.

There is a 35% probability that the 2005 Northwest Pacific typhoon season ACE index will be above average (defined as an ACE index value in the upper tercile historically (>331)), a 51% likelihood it will be near-normal (defined as an ACE index value in the middle tercile historically (237 to 331) and only a 14% chance it will be below-normal (defined as an ACE index value in the lower tercile historically (<237)). The 40-year period 1965-2004 is used for climatology.

Key: Terciles = Data groupings of equal (33.3%) probability corresponding to the upper, middle and lower one third of values historically (1965-2004).

Key Predictors for 2005

The TSR predictors have changed from recent years. This follows a review of seasonal predictability over the extended 1965-2004 period of reliable data. Tropical storm and typhoon numbers are now forecast before May using the Niño 3 sea surface temperature (SST) from the prior September; from May they are forecast using April surface pressure over the region 17.5°N-35°N, 160°E-175°W. Intense typhoon numbers and the ACE index are now forecast in March and April using the February surface pressure in the central northern tropical Pacific region 10°N-20°N, 145°W-165°W; from May they are forecast from the forecast value for the August-September Niño 3.75 index (5°S-5°N, 140°W-180°W). Above average (below average) Niño 3.75 SSTs are associated with weaker (stronger) trade winds over the region 2.5°N-12.5°N, 120°E-180°E. These in turn lead to enhanced (reduced) cyclonic vorticity over the Northwest Pacific region where intense typhoons form. The TSR forecast anomaly (1965-2004) climatology) for August-September Niño 3.75 is $0.28 \pm 0.20^{\circ}\text{C}$ (down from $0.33 \pm 0.30^{\circ}\text{C}$ last month). The forecast skill (1965-2004) for this predictor at this lead is 92%.

Further Information

Further information about the TSR forecasts, verifications and hindcast skill as a function of lead time may be obtained from the TSR website (<http://tropicalstormrisk.com>). A summary of the 2005 Northwest Pacific typhoon season and a verification of the TSR seasonal forecasts will be issued in January 2006.

Appendix - Predictions from Previous Months

a) Deterministic forecasts

NW Pacific ACE Index and System Numbers 2005					
		ACE Index ($\times 10^4$ knots ²)	Tropical Storms	Typhoons	Intense Typhoons
Average Number (\pm SD) (1965-2004)		305 (\pm 99)	26.7 (\pm 4.5)	16.9 (\pm 3.7)	8.6 (\pm 3.0)
TSR Forecasts (\pm FE)	5th August 2005	328 (\pm 78)	27.6 (\pm 3.7)	17.5 (\pm 2.9)	9.4 (\pm 2.5)
	7th July 2005	333 (\pm 83)	27.6 (\pm 3.7)	17.5 (\pm 2.9)	9.5 (\pm 2.5)
	7th June 2005	328 (\pm 84)	27.6 (\pm 3.7)	17.5 (\pm 2.9)	9.4 (\pm 2.7)
	5th May 2005	314 (\pm 80)	27.6 (\pm 3.7)	17.5 (\pm 2.9)	8.9 (\pm 2.6)
	7th March 2005	340 (\pm 92)	25.9 (\pm 4.0)	16.1 (\pm 3.3)	9.8 (\pm 2.7)
Chan Forecasts	24th June 2005	-	25	16	-
	27th April 2005	-	24	15	-

b) Probabilistic forecasts

NW Pacific Total ACE Index 2005				
		Tercile Probabilities		
		below normal	normal	above normal
Climatology 1965-2004		33.3	33.3	33.3
TSR Forecasts	5th August 2005	14	51	35
	7th July 2005	14	48	38
	7th June 2005	16	48	36
	5th May 2005	19	51	30
	7th March 2005	15	43	42

